

ON CONSECUTIVE QUADRATIC NON-RESIDUES: A CONJECTURE OF ISSAI SCHUR

PATRICK HUMMEL

Issai Schur once asked if it was possible to determine a bound, preferably using elementary methods, such that for all prime numbers p greater than the bound, the greatest possible number of consecutive quadratic non-residues modulo p is always less than $p^{1/2}$. (One can find a brief discussion of this problem in R. K. Guy's book [5]). Schur also pointed out that the greatest number of consecutive quadratic non-residues exceeds $p^{1/2}$ for $p = 13$, since 5, 6, 7, and 8 are all quadratic non-residues (mod p). This paper uses elementary methods to prove the following:

Theorem. *$p = 13$ is the only prime number for which the greatest number of consecutive quadratic non-residues modulo p exceeds $p^{1/2}$.*

This problem has been attacked previously using both analytic and elementary methods. We shall briefly consider the results given to us by analytic number theory, and then focus on the elementary methods for the remainder of the paper.

In [4], D. A. Burgess proves the following:

Theorem (D. A. Burgess [4]). *If χ is any non-trivial Dirichlet character of prime modulus p and $\chi(N+1) = \chi(N+2) = \dots = \chi(N+H)$, then $H = O(p^{1/4} \log p)$.*

From this, it follows that there must be some M , such that for all $p > M$, the greatest number of consecutive quadratic non-residues modulo p is less than $p^{1/2}$. In [9], K. K. Norton asserts that he can refine Burgess's method to obtain the following result:

Theorem (K. K. Norton [9]). *In Burgess's theorem, $H < 4.1p^{1/4} \log p$ for all p . If $p > e^{15} \approx 3.27 \times 10^6$, then $H < 2.5p^{1/4} \log p$.*

This result implies that $M = e^{15} \approx 3.27 \times 10^6$ is a suitable value for the aforementioned constant. Unfortunately, however, Norton does not prove this result in his paper, and without a value such as 4.1 for the implied constant in Burgess's theorem, we cannot use Burgess's theorem to find a suitable constant, M , in Schur's conjecture.

Now we consider the work that has been done on the problem using elementary methods. A. Brauer [3] has proved the following theorem:

Theorem (A. Brauer [3]). *For prime numbers p , of the form $4n-1$, the maximum length l , of sequences of quadratic residues and non-residues satisfies $l < p^{1/2}$.*

R. H. Hudson then considers the case $p \equiv 1 \pmod{4}$ by breaking it up into several cases. In [7], he proves that the maximum number of consecutive quadratic non-residues (mod p) is less than $p^{1/2}$ if $p \equiv 1 \pmod{24}$. In [6], he demonstrates that this also holds if $p \equiv 5$ or $p \equiv 17 \pmod{24}$. Putting these together, Hudson obtains the following beneficial result:

Theorem (R. H. Hudson [6]). *If p is a prime, and the greatest number of consecutive quadratic non-residues modulo p exceeds $p^{1/2}$, then $p \equiv 13 \pmod{24}$.*

In the same paper, Hudson proposes a proof that the greatest number of consecutive quadratic non-residues modulo p , is less than $p^{1/2}$ for $p > 2^{332}$ and $p \equiv 13 \pmod{24}$. If his proof of this result were correct, it would complete an elementary proof of Schur's conjecture. But upon a careful reading, one sees that the proof of his assertion is flawed. In particular, Hudson claims that the existence of a quadratic non-residue in the interval $(\frac{p^{1/2}}{128} - 2^{3/2}p^{1/8}, \frac{p^{1/2}}{128})$ implies a quadratic *non-residue* is contained in the interval $(p^{1/2} - 2^{17/2}p^{1/8}, p^{1/2})$, whereas it really implies a quadratic *residue* is contained in the interval $(p^{1/2} - 2^{17/2}p^{1/8}, p^{1/2})$, since 2 is a quadratic non-residue modulo p if $p \equiv 13 \pmod{24}$.

The research presented here tackles the case $p \equiv 13 \pmod{24}$ by using an alternate method. In addition to completing an elementary proof of Schur's conjecture, this paper also shows that $p = 13$ is the only prime number for which the greatest number of consecutive quadratic non-residues $(\bmod p)$ exceeds $p^{1/2}$.

The argument in this paper breaks down into two parts. In the first part, it is shown that if $p \equiv 13 \pmod{24}$ is sufficiently large and if there exists an interval containing more than $p^{1/2}$ integers, all of which are quadratic non-residues modulo p , then there must exist such an interval J satisfying

$$J \subset (\frac{p+3+p^{1/2}}{2}, \frac{p}{2} + 2^{1/2}p^{3/4} - p^{1/2}).$$

Then, in the second part, it is shown that this cannot hold for p sufficiently large by demonstrating that one can find two integers, $a, b \in J$ such that if $R \equiv ab \pmod{p}$ and $0 \leq R \leq p-1$, then $R \in J$. This implies that J contains a quadratic residue R , so J could not have existed in the first place. This implies that the greatest number of consecutive quadratic non-residues modulo p , is less than $p^{1/2}$ when p is sufficiently large. The remaining cases, where p is less than a given bound, are handled by computer.

This research is the result of work done under the mentorship of Prof. Dinakar Ramakrishnan while on a Caltech Summer Undergraduate Research Fellowship. I am deeply grateful to Prof. Ramakrishnan for his guidance and to David Whitehouse for reviewing my paper.

Lemma 1. *If p is a prime number such that $p \equiv 13 \pmod{24}$, $p > 38659$, and there is a sequence of more than $p^{1/2}$ consecutive quadratic non-residues $(\bmod p)$, there must be such a sequence in the interval $(\frac{p+3+p^{1/2}}{2}, \frac{p}{2} + 2^{1/2}p^{3/4} - p^{1/2})$.*

Proof. $p \equiv 13 \pmod{24}$ implies that every number of the form $2a^2$ is a quadratic non-residue. This means that there must exist a quadratic non-residue, say N , in the interval

$$(p^{1/2} - 2^{3/2}p^{1/4} + 2, p^{1/2}),$$

because if c is the smallest positive integer such that $2c^2 > p^{1/2}$, then $2(c-1)^2 > p^{1/2} - 2^{3/2}p^{1/4} + 2$.

Suppose J is an integer interval containing more than $p^{1/2}$ consecutive quadratic non-residues. Multiplying each member of J by N and reducing $(\bmod p)$, we obtain

a collection of quadratic residues in which each quadratic residue differs from the next by $N < p^{1/2}$. This collection must span more than

$$(p^{1/2} - 1)(p^{1/2} - 2^{3/2}p^{1/4} + 2) > p - 2^{3/2}p^{3/4} + p^{1/2}$$

integers. Since -1 is a quadratic residue (mod p), $p - b$ must be a quadratic non-residue whenever b is. Therefore, if our collection of quadratic residues is to lie entirely outside a sequence of more than $p^{1/2}$ consecutive quadratic non-residues, J must either be contained in

$$(1, 2^{1/2}p^{3/4} - \frac{p^{1/2}}{2}),$$

$$(p - 2^{1/2}p^{3/4} + \frac{p^{1/2}}{2}, p - 1),$$

or

$$(1) \quad (\frac{p}{2} - 2^{1/2}p^{3/4} + \frac{p^{1/2}}{2}, \frac{p}{2} + 2^{1/2}p^{3/4} - \frac{p^{1/2}}{2}).$$

But J cannot be fully contained in the first of these intervals because the difference between the square numbers in $(1, 2^{1/2}p^{3/4} - \frac{p^{1/2}}{2})$ is less than $p^{1/2}$. Similarly, J cannot be contained in the second of these intervals because any sequence of the form given by J in $(p - 2^{1/2}p^{3/4} + \frac{p^{1/2}}{2}, p - 1)$ would have to correspond to a similar sequence in $(1, 2^{1/2}p^{3/4} - \frac{p^{1/2}}{2})$. So such a J can only be contained in the interval given by (1).

We now refer to the following theorem of A. Brauer's:

Theorem (A. Brauer [1]). *The least odd quadratic non-residue u modulo a prime p satisfies $u < 2^{3/5}p^{2/5} + 2^{-(6/5)} \cdot 25p^{1/5} + 3$ for $p = 8n + 5$.*

This implies that there exists an odd quadratic non-residue u , less than $p^{1/2}$ if $p > 38659$. Then, since $\frac{p+1}{2}$ is a quadratic non-residue (mod p), $u(\frac{p+1}{2}) \equiv \frac{p+u}{2}$ is a quadratic residue (mod p). Therefore, there exists a quadratic residue in the interval $(\frac{p}{2}, \frac{p+p^{1/2}}{2})$, so there must exist a corresponding quadratic residue in the interval $(\frac{p-p^{1/2}}{2}, \frac{p}{2})$, which means that J cannot pass through $\frac{p}{2}$.

Combining this with the fact that $-b$ is a quadratic non-residue whenever b is, we know that if such a J exists, there must be at least one such J in the interval

$$(2) \quad (\frac{p}{2}, \frac{p}{2} + 2^{1/2}p^{3/4} - \frac{p^{1/2}}{2}).$$

Now note that 3 is a quadratic residue (mod p). Therefore, for odd m , $\frac{p+3m}{2}$ must be a quadratic residue (mod p) if $\frac{p+m}{2}$ is. Combined with the fact that there exists a quadratic residue in the interval $(\frac{p}{2}, \frac{p+p^{1/2}}{2})$, we find that if such a J lies in the interval given by (2), that same J must also lie in the interval

$$(3) \quad (\frac{p+3+p^{1/2}}{2}, \frac{p}{2} + 2^{1/2}p^{3/4} - \frac{p^{1/2}}{2}).$$

To see why, suppose that $\frac{p+3}{2} + x$ is the first entry in J . Then, we can assume that $\frac{p+1}{2} + x$ is a quadratic residue, meaning $\frac{p}{2} + 3(\frac{1}{2} + x) = \frac{p+3}{2} + 3x$ is also a quadratic residue. Therefore we must have $\frac{p+3}{2} + 3x - (\frac{p+3}{2} + x) > p^{1/2}$, or $x > \frac{p^{1/2}}{2}$. $\square$

Lemma 2. *Suppose $p > 38659$ is a prime congruent to 13 modulo 24 and $\frac{p+1}{2} + k$ is a quadratic non-residue, where $k > 0$ is some fixed integer. Then, if there exists a such that $\frac{1}{4} \leq a \leq \frac{15}{32}$, and $(ap^{1/2} - 2)^2 > 2k + 2(1-a)p^{1/2} + 2 - \lfloor p^{1/2} \rfloor$, and the difference between $(\frac{p+1}{2})^2 + k + k^2 + 2kx + x + x^2$ at $x = \lfloor (1-a)p^{1/2} \rfloor$ and $x = \lfloor ap^{1/2} \rfloor - 2$ is greater than p , $\frac{p+1}{2} + k$ is not the smallest number in a sequence of more than $p^{1/2}$ consecutive quadratic non-residues.*

Proof. Suppose that all the integers of the form

$$\frac{p+1}{2} + k + m$$

are quadratic non-residues, where m is an integer ranging from 0 to $\lfloor p^{1/2} \rfloor$.

Note that the product of two integers of this form, say $\frac{p+1}{2} + k + m$ and $\frac{p+1}{2} + k + n$, is a quadratic residue, and equals

$$(4) \quad \left(\frac{p+1}{2}\right)^2 + (p+1)k + k^2 + (m+n)\left(k + \frac{p+1}{2}\right) + mn$$

which if m and n both equal the same value, say x , reduces to

$$\left(\frac{p+1}{2}\right)^2 + (p+1)k + k^2 + (2x)\left(k + \frac{p+1}{2}\right) + x^2$$

which is congruent to

$$(5) \quad \left(\frac{p+1}{2}\right)^2 + k + k^2 + 2kx + x + x^2$$

modulo p .

If there exists an a such that $\frac{1}{4} \leq a \leq \frac{15}{32}$ and the difference between $(\frac{p+1}{2})^2 + k + k^2 + 2kx + x + x^2$ at $x = \lfloor (1-a)p^{1/2} \rfloor$ and $x = \lfloor ap^{1/2} \rfloor - 2$ is greater than p , we note that we can find an integer x contained in the interval $(ap^{1/2} - 2, (1-a)p^{1/2}]$, and an integer c such that

$$\left(\frac{p+1}{2}\right)^2 + k + k^2 + 2kx + x + x^2 > (c + \frac{1}{2})p + \frac{1}{2} + k + \lfloor p^{1/2} \rfloor$$

and

$$(6) \quad \left(\frac{p+1}{2}\right)^2 + k + k^2 + 2k(x-1) + x - 1 + (x-1)^2 \leq (c + \frac{1}{2})p + \frac{1}{2} + k + \lfloor p^{1/2} \rfloor.$$

Now suppose that

$$\left(\frac{p+1}{2}\right)^2 + k + k^2 + 2k(x-1) + x - 1 + (x-1)^2 \geq (c + \frac{1}{2})p + \frac{1}{2} + k.$$

Then, combining this with (6), we reach the absurd conclusion that a quadratic residue equals a quadratic non-residue. Therefore, we have

$$(7) \quad \left(\frac{p+1}{2}\right)^2 + k + k^2 + 2k(x-1) + x - 1 + (x-1)^2 < (c + \frac{1}{2})p + \frac{1}{2} + k.$$

Now consider (4) again. Let m and n vary so that $m = x - y$ and $n = x + y$, where x is an integer that satisfies the above conditions, and y is an integer ranging from 0 to the smallest integer larger than $ap^{1/2} - 2$. Since x lies in $(ap^{1/2} - 2, (1-a)p^{1/2}]$, we continue to meet the condition that m and n are both integers between 0 and $\lfloor p^{1/2} \rfloor$ inclusive, because $y < ap^{1/2} - 1$, and $\lfloor p^{1/2} \rfloor - (1-a)p^{1/2} > ap^{1/2} - 1$. If m and n vary this way, the only part of (4) that changes is the product mn . Also note that

$$0 < mn - (m-1)(n+1) = n - m + 1 < p^{1/2}$$

when $0 \leq n - m < p^{1/2} - 1$, which holds when y varies as above. So we have a collection of quadratic residues in which no quadratic residue exceeds the next by more than $p^{1/2}$. This collection spans an interval of

$$(8) \quad x^2 - (x - ap^{1/2} + 2)(x + ap^{1/2} - 2) = (ap^{1/2} - 2)^2.$$

Note that increasing x by 1 in (5) increases the value of the expression by $2k + 2x + 2$. Combining this with (7) and (8), we find that one of the quadratic residues in the aforementioned collection is congruent (mod p) to an integer in the interval, $(\frac{p+1}{2} + k, \frac{p+1}{2} + k + \lfloor p^{1/2} \rfloor)$ if

$$(ap^{1/2} - 2)^2 > 2k + 2x + 2 - \lfloor p^{1/2} \rfloor,$$

and since $x \leq (1-a)p^{1/2}$, we have

$$(9) \quad (ap^{1/2} - 2)^2 > 2k + 2(1-a)p^{1/2} + 2 - \lfloor p^{1/2} \rfloor,$$

which proves the lemma. □

Theorem. $p = 13$ is the only prime number for which the greatest number of consecutive quadratic non-residues modulo p exceeds $p^{1/2}$.

Proof. Suppose $p > 38659$, and suppose there exists a sequence of more than $p^{1/2}$ consecutive quadratic non-residues (mod p). As noted earlier, this implies that $p \equiv 13 \pmod{24}$.

Now suppose that

$$(10) \quad \frac{p+1}{2} + k$$

is a quadratic non-residue, where k is a fixed integer. With Lemma 1 in mind, we need only prove that this is not the least quadratic non-residue in a sequence of more than $p^{1/2}$ consecutive quadratic non-residues when

$$\frac{p^{1/2}}{2} + 1 < k < 2^{1/2}p^{3/4} - p^{1/2}.$$

Now consider three cases:

Case 1: $k < 2p^{1/2}$. Note that the difference between (5) at $x = \lfloor \frac{3p^{1/2}}{4} \rfloor$ and $x = \lfloor \frac{p^{1/2}}{4} \rfloor - 2$ is greater than p because it equals

$$\begin{aligned} & \left((2k+1) \left\lfloor \frac{3p^{1/2}}{4} \right\rfloor + \left\lfloor \frac{3p^{1/2}}{4} \right\rfloor^2 \right) - \left((2k+1) \left(\left\lfloor \frac{p^{1/2}}{4} \right\rfloor - 2 \right) + \left(\left\lfloor \frac{p^{1/2}}{4} \right\rfloor - 2 \right)^2 \right) \\ &= 2k \left(\left\lfloor \frac{3p^{1/2}}{4} \right\rfloor - \left\lfloor \frac{p^{1/2}}{4} \right\rfloor \right) + \left\lfloor \frac{3p^{1/2}}{4} \right\rfloor + 4k + \left\lfloor \frac{3p^{1/2}}{4} \right\rfloor^2 - \left\lfloor \frac{p^{1/2}}{4} \right\rfloor^2 + 3 \left\lfloor \frac{p^{1/2}}{4} \right\rfloor - 2 \\ &> 2k \left(\frac{p^{1/2}}{2} - 1 \right) + \frac{3p^{1/2}}{4} - 1 + 4k + \frac{9p}{16} - \frac{3p^{1/2}}{2} + 1 - \frac{p}{16} + 3 \left(\frac{p^{1/2}}{4} - 1 \right) - 2 \\ &= kp^{1/2} + 2k + \frac{p}{2} - 5 > \frac{p}{2} + p^{1/2} + p^{1/2} + 2 + \frac{p}{2} - 5 > p, \end{aligned}$$

since $k > \frac{p^{1/2}}{2} + 1$. With Lemma 2 in mind, we find that (10) is not the least quadratic non-residue in a sequence of more than $p^{1/2}$ consecutive quadratic non-residues if

$$(11) \quad \left(\frac{p^{1/2}}{4} - 2 \right)^2 > 2k + \frac{3p^{1/2}}{2} + 2 - \lfloor p^{1/2} \rfloor.$$

Since $k < 2p^{1/2}$, (11) holds whenever

$$\left(\frac{p^{1/2}}{4} - 2 \right)^2 > \frac{9p^{1/2}}{2} + 3,$$

which holds for $p > 7711$.

Case 2: $2p^{1/2} < k < 8p^{1/2}$. Note that the difference between (5) at $x = \lfloor \frac{5p^{1/2}}{8} \rfloor$ and $x = \lfloor \frac{3p^{1/2}}{8} \rfloor - 2$ is greater than p because it equals

$$\begin{aligned} & \left((2k+1) \left\lfloor \frac{5p^{1/2}}{8} \right\rfloor + \left\lfloor \frac{5p^{1/2}}{8} \right\rfloor^2 \right) - \left((2k+1) \left(\left\lfloor \frac{3p^{1/2}}{8} \right\rfloor - 2 \right) + \left(\left\lfloor \frac{3p^{1/2}}{8} \right\rfloor - 2 \right)^2 \right) \\ &= 2k \left(\left\lfloor \frac{5p^{1/2}}{8} \right\rfloor - \left\lfloor \frac{3p^{1/2}}{8} \right\rfloor \right) + \left\lfloor \frac{5p^{1/2}}{8} \right\rfloor + 4k + \left\lfloor \frac{5p^{1/2}}{8} \right\rfloor^2 - \left\lfloor \frac{3p^{1/2}}{8} \right\rfloor^2 + 3 \left\lfloor \frac{3p^{1/2}}{8} \right\rfloor - 2 \\ &> 2k \left(\frac{p^{1/2}}{4} - 1 \right) + \frac{5p^{1/2}}{8} - 1 + 4k + \frac{25p}{64} - \frac{5p^{1/2}}{4} + 1 - \frac{9p}{64} + 3 \left(\frac{3p^{1/2}}{8} - 1 \right) - 2 \\ &= \frac{kp^{1/2}}{2} + 2k + \frac{p}{4} + \frac{p^{1/2}}{2} - 5 > p + 4p^{1/2} + \frac{p}{4} + \frac{p^{1/2}}{2} - 5 > p, \end{aligned}$$

since $k > 2p^{1/2}$. With Lemma 2 in mind, we find that (10) is not the least quadratic non-residue in a sequence of more than $p^{1/2}$ consecutive quadratic non-residues if

$$(12) \quad \left(\frac{3p^{1/2}}{8} - 2 \right)^2 > 2k + \frac{5p^{1/2}}{4} + 2 - \lfloor p^{1/2} \rfloor.$$

Since $k < 8p^{1/2}$, (12) holds when

$$\left(\frac{3p^{1/2}}{8} - 2\right)^2 > \frac{65p^{1/2}}{4} + 3,$$

which holds for $p > 15917$.

Case 3: $8p^{1/2} < k$. Note that the difference between (5) at $x = \lfloor \frac{17p^{1/2}}{32} \rfloor$ and $x = \lfloor \frac{15p^{1/2}}{32} \rfloor - 2$ is greater than p because it equals

$$\begin{aligned} & \left((2k+1) \left\lfloor \frac{17p^{1/2}}{32} \right\rfloor + \left\lfloor \frac{17p^{1/2}}{32} \right\rfloor^2 \right) - \left((2k+1) \left(\left\lfloor \frac{15p^{1/2}}{32} \right\rfloor - 2 \right) + \left(\left\lfloor \frac{15p^{1/2}}{32} \right\rfloor - 2 \right)^2 \right) \\ &= 2k \left(\left\lfloor \frac{17p^{1/2}}{32} \right\rfloor - \left\lfloor \frac{15p^{1/2}}{32} \right\rfloor \right) + \left\lfloor \frac{17p^{1/2}}{32} \right\rfloor^2 + 4k + \left\lfloor \frac{17p^{1/2}}{32} \right\rfloor - \left\lfloor \frac{15p^{1/2}}{32} \right\rfloor^2 + 3 \left\lfloor \frac{15p^{1/2}}{32} \right\rfloor - 2 \\ &> 2k \left(\frac{p^{1/2}}{16} - 1 \right) + \frac{17p^{1/2}}{32} - 1 + 4k + \frac{289p}{1024} - \frac{17p^{1/2}}{16} + 1 - \frac{225p}{1024} + 3 \left(\frac{15p^{1/2}}{32} - 1 \right) - 2 \\ &= \frac{kp^{1/2}}{8} + 2k + \frac{p}{16} + \frac{7p^{1/2}}{8} - 5 > p + 16p^{1/2} + \frac{p}{16} + \frac{7p^{1/2}}{8} - 5 > p, \end{aligned}$$

since $k > 8p^{1/2}$. With Lemma 2 in mind, we find that (10) is not the least quadratic non-residue in a sequence of more than $p^{1/2}$ consecutive quadratic non-residues if

$$(13) \quad \left(\frac{15p^{1/2}}{32} - 2 \right)^2 > 2k + \frac{17p^{1/2}}{16} + 2 - \lfloor p^{1/2} \rfloor.$$

Since $k < 2^{1/2}p^{3/4} - p^{1/2}$, (13) holds whenever

$$\left(\frac{15p^{1/2}}{32} - 2 \right)^2 > 2^{3/2}p^{3/4} - \frac{31p^{1/2}}{16} + 3,$$

which holds for $p > 27250$.

So when $p > 38659$, no sequence of more than $p^{1/2}$ consecutive quadratic non-residues exists.

Now all that remains is to consider the case $p \leq 38659$. This case can be handled by a simple computation. I have run a computer program which compares the largest number of consecutive quadratic non-residues modulo p with $p^{1/2}$ for all primes p , such that $p \equiv 13 \pmod{24}$ and $p \leq 38659$. From this I was able to check that 13 is the only prime number for which the greatest number of consecutive quadratic non-residues (mod p) exceeds $p^{1/2}$.

□

Remark. The data obtained from this program can be viewed by going to the website <http://www.math.caltech.edu/people/hummel.html>. A sample of some of the data obtained from the program is given below. The numbers in each set represent p , the greatest number of consecutive quadratic non-residues (mod p), and $\lfloor p^{1/2} \rfloor$ in that order. For all but the smallest numbers, $p^{1/2}$ far exceeds the greatest number of consecutive quadratic non-residues.

{13, 4, 3}, {757, 8, 27}, {3181, 9, 56}, {5869, 9, 76}, {7237, 10, 85}, {9397, 10, 96}, {12037, 11, 109}, {14389, 12, 119}, {16477, 12, 128}, {18517, 13, 136}, {20509, 13, 143},

$\{22381, 12, 149\}, \{24061, 13, 155\}, \{26029, 13, 161\}, \{28429, 13, 168\}, \{30469, 14, 174\},$
 $\{32749, 15, 180\}, \{34693, 14, 186\}, \{36709, 15, 191\}, \{38653, 15, 196\}.$

REFERENCES

- [1] A. Brauer, On the non-existence of the Euclidean algorithm in certain quadratic number fields, *Amer. J. Math.* **62** (1940), 697-716.
- [2] A. Brauer, Über den kleinsten quadratischen Nichtrest, *Math. Zeitschr.* **33** (1930), 161-176.
- [3] A. Brauer, Über die Verteilung der Potenzreste, *Math. Zeitschr.* **35** (1932), 39-50.
- [4] D. A. Burgess, A note on the distribution of residues and non-residues, *J. London Math. Soc.* **38** (1963), 253-256.
- [5] R. K. Guy, "Unsolved Problems in Number Theory," Springer-Verlag, New York (1994), 244-245.
- [6] R. H. Hudson, On a conjecture of Issai Schur, *J. Reine Angew. Math.* **289** (1977), 215-220.
- [7] R. H. Hudson, On sequences of consecutive quadratic non-residues, *J. Number Theory* **3** (1971), 178-181.
- [8] R. H. Hudson, Prime k-th power non-residues, *Acta Arithmetica* **23** (1973), 89-106.
- [9] K. K. Norton, Bounds for sequences of consecutive power residues, *Analytic number theory*, Amer. Math. Soc. **24** (1973), 213-220.

Patrick Hummel is affiliated with the Class of 2006 at California Institute of Technology.

1170 E. DEL MAR BLVD., APT. 10 PASADENA, CA 91106-3425
E-mail address: hummel@its.caltech.edu